

Important Safety Information (Continued)

MYOBLOC is contraindicated in patients with a known hypersensitivity to any botulinum toxin preparation or to any of the components in the formulation.

MYOBLOC is contraindicated for use in patients with infection at the proposed injection site(s).

The potency Units of MYOBLOC are specific to the preparation and assay method utilized. They are not interchangeable with other preparations of botulinum toxin products and, therefore, units of biological activity of MYOBLOC cannot be compared to or converted into units of any other botulinum toxin products assessed with any other specific assay method.

Treatment with MYOBLOC and other botulinum toxin products can result in swallowing or breathing difficulties. Patients with pre-existing swallowing or breathing difficulties may be more susceptible to these complications. In most cases, this is a consequence of weakening of muscles in the area of injection that are involved in breathing or swallowing. When distant effects occur, additional respiratory muscles may be involved.

Deaths as a complication of severe dysphagia have been reported after treatment with botulinum toxin. Dysphagia may persist for several months, and require use of a feeding tube to maintain adequate nutrition and hydration. Aspiration may result from severe dysphagia and is a particular risk when treating patients in whom swallowing or respiratory function is already compromised.

Treatment of cervical dystonia with botulinum toxins may weaken neck muscles that serve as accessory muscles of ventilation. This may result in a critical loss of breathing capacity in patients with respiratory disorders who may have become dependent upon these accessory muscles. There have been postmarketing reports of serious breathing difficulties, including respiratory failure, in cervical dystonia patients. Patients treated with botulinum toxin may require immediate medical attention should they develop problems with swallowing, speech or respiratory disorders. These reactions can occur within hours to weeks after injection with botulinum toxin.

Individuals with peripheral motor neuropathic diseases, amyotrophic lateral sclerosis, or neuromuscular junctional disorders (e.g., myasthenia gravis or Lambert-Eaton syndrome) should be monitored particularly closely when given botulinum toxin. Patients with neuromuscular disorders may be at increased risk of clinically significant effects including severe dysphagia and respiratory compromise from typical doses of MYOBLOC.

This product contains albumin, a derivative of human blood. Based on effective donor screening and product manufacturing processes, it carries an extremely remote risk for transmission of viral diseases. A theoretical risk for transmission of Creutzfeldt-Jakob disease (CJD) also is considered extremely remote. No cases of transmission of viral diseases or CJD have ever been identified for albumin.

Only 9 subjects without a prior history of tolerating injections of type A botulinum toxin have been studied. Treatment of botulinum toxin naïve patients should be initiated at lower doses of MYOBLOC.

Co-administration of MYOBLOC and aminoglycosides or other agents interfering with neuromuscular transmission (e.g., curare-like compounds) should only be performed with caution as the effect of the toxin may be potentiated.

The effect of administering different botulinum neurotoxin serotypes at the same time or within less than 4 months of each other is unknown. However, neuromuscular paralysis may be potentiated by co-administration or overlapping administration of different botulinum toxin serotypes.

It is not known whether MYOBLOC can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. MYOBLOC should be given to a pregnant woman only if clearly needed.

The most commonly reported adverse events associated with MYOBLOC treatment in all studies were dry mouth, dysphagia, dyspepsia, and injection site pain. Dry mouth and dysphagia were the adverse reactions most frequently resulting in discontinuation of treatment. There was an increased incidence of dysphagia with increased dose in the sternocleidomastoid muscle. The incidence of dry mouth showed some dose-related increase with doses injected into the splenius capitis, trapezius and sternocleidomastoid muscles.

To report SUSPECTED ADVERSE REACTIONS or product complaints, contact US WorldMeds at 1-888-461-2255, Option 2. You may also report SUSPECTED ADVERSE REACTIONS to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see accompanying full Prescribing Information, including Boxed WARNING, and Medication Guide.

SOLSTICE[®]
NEUROSCIENCES
A US WORLDMEDS COMPANY

MYOBLOC[®]
rimabotulinumtoxinB
Injection [5,000 Units/mL]
www.myobloc.com

US WorldMeds

MYOBLOC[®]
rimabotulinumtoxinB
Injection [5,000 Units/mL]

Receive up to \$4,000
each year to help with your
out-of-pocket expenses for MYOBLOC

ENROLL IN THE MYOBLOC COPAY PROGRAM TODAY!

No Financial Requirements

Indication

MYOBLOC (rimabotulinumtoxinB) Injection is indicated for the treatment of adults with cervical dystonia to reduce the severity of abnormal head position and neck pain associated with cervical dystonia.

Important Safety Information

WARNING: DISTANT SPREAD OF TOXIN EFFECT

Postmarketing reports indicate that the effects of MYOBLOC and all botulinum toxin products may spread from the area of injection to produce symptoms consistent with botulinum toxin effects. These may include asthenia, generalized muscle weakness, diplopia, blurred vision, ptosis, dysphagia, dysphonia, dysarthria, urinary incontinence, and breathing difficulties. These symptoms have been reported hours to weeks after injection. Swallowing and breathing difficulties can be life threatening and there have been reports of death. The risk of symptoms is probably greatest in children treated for spasticity but symptoms can also occur in adults treated for spasticity and other conditions, particularly in those patients who have underlying conditions that would predispose them to these symptoms. In unapproved uses, including spasticity in children and adults, and in approved indications, cases of spread of effect have occurred at doses comparable to those used to treat cervical dystonia and at lower doses [see Warnings and Precautions].

Please see the full Important Safety Information on back page and the full Prescribing Information, including Boxed WARNING, and Medication Guide in inside pocket.

MYOBLOC COPAY PROGRAM



WHO'S ELIGIBLE

Patients who:

- Are 18 years or older and a legal U.S. resident
- Have commercial insurance coverage
- Are diagnosed with cervical dystonia (G24.3)
- Are NOT enrolled in a government insurance plan (e.g., Medicare, Medicaid, TRICARE[®], and other federal- or state-funded programs)



WHAT'S COVERED

- Any size vial of MYOBLOC
- Injection procedure costs
- Injection guidance costs
 - Ultrasound or EMG
- Residents of Michigan, Rhode Island, and Minnesota are not eligible for injection procedure or injection guidance costs



IT'S EASY TO ENROLL

**Your healthcare provider can enroll you
OR you can call 1.877.268.7697 to enroll.**

- No enrollment forms needed
- No financial requirements

Please see the full Important Safety Information on back page and the full Prescribing Information, including Boxed WARNING, and Medication Guide in inside pocket.



HOW MUCH CAN I SAVE

- Up to \$4,000 per year
- There are NO cost limitations per injection



ENROLLMENT COMPLETE. NOW WHAT?

Once enrollment is completed and your Explanation of Benefits (EOB) from your insurance provider is received:

Your out-of-pocket expenses can be paid

1. Directly to your site of care on your behalf (administering office, specialty pharmacy, etc.)
2. To you as a reimbursement for out-of-pocket expenses you paid to the site of care

**Start Saving Today on your MYOBLOC Treatments
Call 1.877.268.7697**

Terms and Conditions for Patients: 1. This offer is valid for commercially-insured patients only and is good for use only with a MYOBLOC prescription at the time the prescription is filled or after the product is administered to the patient. 2. Depending on insurance coverage, eligible, insured patients may pay no more than zero dollars (\$0) for MYOBLOC and the administrative services associated with MYOBLOC, up to a maximum savings limit of four thousand dollars (\$4,000) per year. Patient out-of-pocket expense may vary. 3. This offer is not valid for patients enrolled in Medicare, Medicaid, or other federal or state healthcare programs, or private indemnity or HMO insurance plans that reimburse you for the entire cost of your prescription drugs. Patients may not use this Program if they are Medicare-eligible and enrolled in an employer-sponsored health plan or medical or prescription drug benefit program for retirees. 4. The offer is valid for one (1) year. 5. US WorldMeds reserves the right to rescind, revoke, or amend this offer without notice. 6. Offer good only in the USA, including Puerto Rico, at participating pharmacies or healthcare providers. 7. Void if prohibited by law, taxed, or restricted. 8. Residents of Michigan, Rhode Island, and Minnesota are not eligible for assistance with payment for injection or injection guidance-related costs, but may receive assistance with MYOBLOC. 9. This Program is not transferable. The selling, purchasing, trading, or counterfeiting of this Program is prohibited by law. 10. This Program is not insurance. 11. By redeeming this assistance, you represent that you are an eligible patient and that you understand and agree to comply with the terms and conditions of this offer.