

## DO YOU EXPERIENCE EXCESSIVE DROOLING?

YOU MAY HAVE CHRONIC SIALORRHEA.

Print out this questionnaire, fill out the answers,  
and share with your doctor to see if MYOBLOC is right for you.

### What is MYOBLOC?

MYOBLOC® is a prescription medicine used in adults that is injected into glands that make saliva and is used to treat long-lasting (chronic) drooling (sialorrhea).

It is not known whether MYOBLOC is safe or effective in children.

### IMPORTANT SAFETY INFORMATION

#### What is the most important information I should know about MYOBLOC?

**MYOBLOC may cause serious side effects that can be life threatening. Call your doctor or get medical help right away if you have any of these problems after treatment with MYOBLOC:**

- **Problems swallowing, speaking, or breathing.** These problems can happen hours to weeks after an injection of MYOBLOC if the muscles that you use to breathe and swallow become weak and death can happen if you have severe problems.
  - People with breathing problems who use muscles in their neck to help them breathe may be at greater risk.
  - Swallowing problems may last several months and require a feeding tube. If severe, food or liquids may go into your lungs. People who already have swallowing or breathing problems are at highest risk.
- **Spread of toxin effects.** In some cases, botulinum toxin may affect areas of the body away from the injection site and cause symptoms of botulism which include: loss of strength and muscle weakness all over the body; trouble saying words clearly (dysarthria); double vision; loss of bladder control; blurred vision and drooping eyelids; trouble breathing; hoarseness or change or loss of voice (dysphonia); trouble swallowing (dysphagia).

**These symptoms can happen hours to weeks after you receive an injection of MYOBLOC.**

These problems could make it unsafe for you to drive a car or do other dangerous activities.

**Please see additional Important Safety Information on [pages 3 and 4](#) and full [Prescribing Information](#) and [Medication Guide](#), including Warning.**

# RECOGNIZE AND RECORD YOUR SYMPTOMS AND EXPERIENCE

---

Symptom recognition often represents the first step toward treatment. Answer the questions below and print out this page to share with your healthcare provider at your next appointment.

- Does saliva dribble in the corner of your mouth during the day?  
 Yes       No
- Does your pillow or bedding get wet from drooling while you sleep?  
 Yes       No
- Does your clothing get wet from drooling during the day?  
 Yes       No
- Do you carry tissues or a handkerchief in case you need to wipe away saliva?  
 Yes       No
- Do you experience irritated skin or infections around the mouth?  
 Yes       No
- Do you experience frequent dehydration?  
 Yes       No
- Do you experience chronic cough?  
 Yes       No
- At times, do you find it difficult to breathe?  
 Yes       No
- At times, do you find it difficult to eat or swallow?  
 Yes       No
- Do you ever feel embarrassed about your drooling?  
 Yes       No

Did you answer yes to any of these questions? MYOBLOC® (rimabotulinumtoxinB) may be right for you. Use this questionnaire to start a conversation with your healthcare provider.

## IMPORTANT SAFETY INFORMATION (cont'd)

### Do not receive MYOBLOC if you:

- are allergic to MYOBLOC or any of the ingredients in MYOBLOC.
- had an allergic reaction to any other botulinum toxin product such as BOTOX®, BOTOX® Cosmetic (onabotulinumtoxinA), DYSPORT® (abobotulinumtoxinA), or XEOMIN® (incobotulinumtoxinA).
- have a skin infection at the planned injection site.

Please see additional Important Safety Information on [pages 3 and 4](#) and full [Prescribing Information and Medication Guide](#), including Warning.

## What is MYOBLOC?

MYOBLOC® is a prescription medicine used in adults that is injected into glands that make saliva and is used to treat long-lasting (chronic) drooling (sialorrhea).

It is not known whether MYOBLOC is safe or effective in children.

## IMPORTANT SAFETY INFORMATION

### What is the most important information I should know about MYOBLOC?

**MYOBLOC may cause serious side effects that can be life threatening. Call your doctor or get medical help right away if you have any of these problems after treatment with MYOBLOC:**

- **Problems swallowing, speaking, or breathing.** These problems can happen hours to weeks after an injection of MYOBLOC if the muscles that you use to breathe and swallow become weak and death can happen if you have severe problems.
  - People with breathing problems who use muscles in their neck to help them breathe may be at greater risk.
  - Swallowing problems may last several months and require a feeding tube. If severe, food or liquids may go into your lungs. People who already have swallowing or breathing problems are at highest risk.
- **Spread of toxin effects.** In some cases, botulinum toxin may affect areas of the body away from the injection site and cause symptoms of botulism which include: loss of strength and muscle weakness all over the body; trouble saying words clearly (dysarthria); double vision; loss of bladder control; blurred vision and drooping eyelids; trouble breathing; hoarseness or change or loss of voice (dysphonia); trouble swallowing (dysphagia).

**These symptoms can happen hours to weeks after you receive an injection of MYOBLOC.**

These problems could make it unsafe for you to drive a car or do other dangerous activities.

### Do not receive MYOBLOC if you:

- are allergic to MYOBLOC or any of the ingredients in MYOBLOC.
- had an allergic reaction to any other botulinum toxin product such as BOTOX®, BOTOX® Cosmetic (onabotulinumtoxinA), DYSPORT® (abobotulinumtoxinA), or XEOMIN® (incobotulinumtoxinA).
- have a skin infection at the planned injection site.

### Before receiving MYOBLOC, tell your doctor about all your medical conditions, including if you have:

a disease that affects your muscles and nerves (such as amyotrophic lateral sclerosis [ALS or Lou Gehrig's disease], myasthenia gravis or Lambert-Eaton syndrome); had any side effect from any botulinum toxin product in the past; a breathing problem, such as asthma or emphysema; a history of swallowing problems or inhaling food or fluid into your lungs (aspiration); bleeding problems; drooping eyelids; plans to have surgery; had surgery on your face.

### Tell your doctor if you:

- are pregnant or plan to become pregnant. It is not known if MYOBLOC can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if MYOBLOC passes into breast milk.

**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins and herbal supplements. **Talk to your doctor before you take any new medicine after you receive MYOBLOC.**

Using MYOBLOC with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you have received MYOBLOC in the past. Especially tell your doctor if you:** have received any other botulinum toxin product in the last 4 months; have received injections of botulinum toxin such as BOTOX®, BOTOX® Cosmetic (onabotulinumtoxinA), DYSPORT® (abobotulinumtoxinA), or XEOMIN® (incobotulinumtoxinA) in the past. Be sure your doctor knows exactly which product you received; have recently received an antibiotic by injection; take muscle relaxants; take an allergy or cold medicine; take a sleep medicine.

### **What should I avoid while receiving MYOBLOC?**

MYOBLOC may cause loss of strength or general muscle weakness or vision problems within hours to weeks of receiving MYOBLOC. **If this happens, do not drive a car, operate machinery or do other dangerous activities.**

### **What are the possible side effects of MYOBLOC?**

**MYOBLOC can cause serious side effects including:**

- allergic reactions. Symptoms of an allergic reaction to MYOBLOC may include: itching, rash, redness, swelling, wheezing, trouble breathing, or dizziness or feeling faint. Tell your doctor or get medical help right away if you get wheezing, or trouble breathing, or if you get dizzy or faint.

**The most common side effects of MYOBLOC in people with sialorrhea include** dry mouth and trouble swallowing (dysphagia).

These are not all the possible side effects of MYOBLOC. **Call your doctor for medical advice about side effects.**

**You are encouraged to report negative side effects of prescription drugs to the FDA.**

**Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.**

**Please see [Medication Guide](#), including Warning, for additional information and talk to your healthcare provider.**

# COULD MYOBLOC BE THE RIGHT TREATMENT FOR YOU?



In clinical studies, MYOBLOC decreased symptoms of chronic sialorrhea in as little as 1 week and can last up to 3 months.

**TALK TO YOUR PHYSICIAN ABOUT GETTING MYOBLOC AND VISIT [www.myobloc.com/For-CS](http://www.myobloc.com/For-CS) TO LEARN MORE ABOUT CHRONIC SIALORRHEA AND HOW MYOBLOC MAY BE ABLE TO HELP.**

## What is MYOBLOC?

MYOBLOC® is a prescription medicine used in adults that is injected into glands that make saliva and is used to treat long-lasting (chronic) drooling (sialorrhea).

It is not known whether MYOBLOC is safe or effective in children.

## IMPORTANT SAFETY INFORMATION

### What is the most important information I should know about MYOBLOC?

**MYOBLOC may cause serious side effects that can be life threatening. Call your doctor or get medical help right away if you have any of these problems after treatment with MYOBLOC:**

- **Problems swallowing, speaking, or breathing.** These problems can happen hours to weeks after an injection of MYOBLOC if the muscles that you use to breathe and swallow become weak and death can happen if you have severe problems.
  - People with breathing problems who use muscles in their neck to help them breathe may be at greater risk.
  - Swallowing problems may last several months and require a feeding tube. If severe, food or liquids may go into your lungs. People who already have swallowing or breathing problems are at highest risk.
- **Spread of toxin effects.** In some cases, botulinum toxin may affect areas of the body away from the injection site and cause symptoms of botulism which include: loss of strength and muscle weakness all over the body; trouble saying words clearly (dysarthria); double vision; loss of bladder control; blurred vision and drooping eyelids; trouble breathing; hoarseness or change or loss of voice (dysphonia); trouble swallowing (dysphagia).

**These symptoms can happen hours to weeks after you receive an injection of MYOBLOC.**

These problems could make it unsafe for you to drive a car or do other dangerous activities.

**Please see additional Important Safety Information on [pages 3 and 4](#) and full [Prescribing Information and Medication Guide](#), including Warning.**

MYOBLOC is a registered trademark of Solstice Neurosciences, LLC, a wholly-owned subsidiary of Supernus Pharmaceuticals, Inc. All other trademarks are the property of their respective owners.  
© 2023. Supernus Pharmaceuticals, Inc. or its related companies. All rights reserved. MYO.2023-0029 05/23.

